

Comprehensive geriatric assessment

Authors: [Katherine T Ward, MD](#), [David B Reuben, MD](#)

Section Editor: [Kenneth E Schmader, MD](#)

Deputy Editor: [Jane Givens, MD](#)

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

Literature review current through: Nov 2018. | **This topic last updated:** Oct 19, 2018.

INTRODUCTION — Geriatric conditions such as functional impairment and dementia are common and frequently unrecognized or inadequately addressed in older adults. Identifying geriatric conditions by performing a geriatric assessment can help clinicians manage these conditions and prevent or delay their complications.

"Geriatric syndrome" is a term that is often used to refer to common health conditions in older adults that do not fit into distinct organ-based disease categories and often have multifactorial causes. The list includes conditions such as cognitive impairment, delirium, incontinence, malnutrition, falls, gait disorders, pressure ulcers, sleep disorders, sensory deficits, fatigue, and dizziness. These conditions are common in older adults, and they may have a major impact on quality of life and disability. Geriatric syndromes can best be identified by a geriatric assessment.

Although the geriatric assessment is a diagnostic process, the term is often used to include both evaluation and management. Geriatric assessment is sometimes used to refer to evaluation by the individual clinician (usually a primary care clinician or a geriatrician) and at other times is used to refer to a more intensive multidisciplinary program, also known as a comprehensive geriatric assessment (CGA).

This topic will review the indications for CGA, as well as its major components and evidence of its efficacy. General issues of geriatric health maintenance and the assessment of specific geriatric populations are discussed elsewhere. (See ["Geriatric health maintenance"](#) and ["Comprehensive geriatric assessment for patients with cancer"](#) and ["Failure to thrive in elderly adults: Evaluation"](#).)

BACKGROUND — Comprehensive geriatric assessment (CGA) is defined as a multidisciplinary diagnostic and treatment process that identifies medical, psychosocial, and functional limitations of a frail older person in order to develop a coordinated plan to maximize overall health with aging [1,2]. The health care of an older adult extends beyond the traditional medical management of illness. It requires evaluation of multiple issues, including physical, cognitive, affective, social, financial, environmental, and spiritual components that influence an older adult's health. CGA is based on the premise that a systematic evaluation of frail, older persons by a team of health professionals may identify a variety of treatable health problems and lead to better health outcomes.

CGA programs are usually initiated through a referral by the primary care clinician or by a clinician caring for a patient in the hospital setting. The content of the assessment varies depending on different settings of care (eg, home, clinic, hospital, nursing home). CGA is not available in all settings, due to issues related to the time required for evaluation, need for coordination of multidisciplinary specialties, and lack of reimbursement for some components (eg, outpatient social work, pharmacy, and nutrition).

INDICATIONS FOR REFERRAL — The best evidence for comprehensive geriatric assessment (CGA) is based on identifying appropriate patients (ie, excluding patients who are either too well or are too sick to

derive benefit). No criteria have been validated to readily identify patients who are likely to benefit from CGA. Specific criteria used by CGA programs to identify patients include:

- Age
- Medical comorbidities such as heart failure or cancer
- Psychosocial disorders such as depression or isolation
- Specific geriatric conditions such as dementia, falls, or functional disability
- Previous or predicted high health care utilization
- Consideration of change in living situation (eg, from independent living to assisted living, nursing home, or in-home caregivers)

One outpatient approach would be to refer patients for CGA who are found to have problems in multiple areas during geriatric assessment screens. Major illnesses (eg, those requiring hospitalization or increased home resources to manage medical and functional needs) should also prompt referral for CGA, particularly for functional status, fall risk, cognitive problems, and mood disorders. (See "[Geriatric health maintenance](#)" and '[Major components](#)' below.)

An inpatient approach would be to refer older patients admitted for a specific medical or surgical reason (eg, fractures, failure to thrive, recurrent pneumonia, pressure sores). Another approach would be to have all patients above a certain age (eg, 85 years) or above a threshold on instruments predicting hospital readmission [3] receive preliminary screening to determine whether a full multidisciplinary evaluation is needed.

Most outpatient CGA programs exclude patients who are unlikely to benefit because of terminal illness, severe dementia, complete functional dependence, and inevitable nursing home placement. However, some of these patients (eg, those with severe dementia) may benefit by increasing the capabilities of caregivers when the assessment is accompanied by ongoing care management. Exclusionary criteria have also included identifying older persons who are "too healthy" to benefit, such as those who are completely functional without any medical comorbidities.

ASSESSMENT TEAM — The range of health care professionals working in the assessment team varies based on the services provided by individual comprehensive geriatric assessment (CGA) programs. In many settings, the CGA process relies on a core team consisting of a clinician, nurse, and social worker and, when appropriate, draws upon an extended team of physical and occupational therapists, nutritionists, pharmacists, psychiatrists, psychologists, dentists, audiologists, podiatrists, and opticians. Although these professionals are usually on-staff in the hospital setting and are also available in the community, access to and reimbursement for these services have limited the availability of CGA programs. Increasingly, CGA programs are moving towards a "virtual team" concept in which members are included as needed, assessments are conducted at different locations on different days, and team communication is completed via telephone or electronically, often through the electronic health record.

Traditionally, the various components of the evaluation are completed by different members of the team, with considerable variability in the assessments. The medical assessment of older persons may be conducted by a physician (usually a geriatrician), nurse practitioner, or physician assistant. The core team (geriatrician, nurse, social worker) may conduct only brief initial assessments or screens for some dimensions. These may be subsequently augmented with more in-depth evaluations by additional professionals. As an example, a dietitian may be needed to assess dietary intake and provide recommendations on optimizing nutrition, or an audiologist may need to conduct a more extensive assessment of hearing loss and evaluate an older person for a hearing aid.

CONDUCTING THE ASSESSMENT

Framework — Conceptually, comprehensive geriatric assessment (CGA) involves several processes of care that are shared over several providers in the assessment team. The overall care rendered by CGA teams can be divided into six steps:

- Data-gathering
- Discussion among the team, increasingly including the patient and/or caregiver as a member of the team
- Development, with the patient and/or caregiver, of a treatment plan
- Implementation of the treatment plan
- Monitoring response to the treatment plan
- Revising the treatment plan

Each of these steps is essential if the process is to be successful at achieving maximal health and functional benefits.

Several different models for CGA have been implemented in various health care settings. Some CGA programs rely on post-discharge assessment due to the decrease in length of hospital stay. Furthermore, while most of the early CGA programs focused on restorative or rehabilitative goals (tertiary prevention), many newer programs are aimed at primary and secondary prevention. (See "[Geriatric health maintenance](#)", [section on 'Overview of prevention for older adults'](#).)

Assessment tools — Although the amount of potentially important information may seem overwhelming, formal assessment tools and shortcuts can reduce this burden on the clinician performing the initial CGA [4]. A pre-visit questionnaire sent to the patient or caregiver prior to the initial assessment can be a timesaving method to gather a large amount of information ([table 1](#)) [5]. These questionnaires can also be completed through secure portals of electronic health records.

These questionnaires can be used to gather information about general history (eg, past medical history, medications, social history, review of systems), as well as gather information specific to CGA, such as:

- Ability to perform functional tasks and need for assistance
- Fall history
- Urinary and/or fecal incontinence
- Pain
- Sources of social support, particularly family or friends
- Depressive symptoms
- Vision or hearing difficulties
- Whether the patient has specified a durable power of attorney for health care

Office staff can be trained to administer screening instruments to both save time and help the clinician to hone in on specific disabilities that need more detailed evaluation ([table 1](#)) [6].

MAJOR COMPONENTS — Core components of comprehensive geriatric assessment (CGA) that should be evaluated during the assessment process are as follows:

- Functional capacity
- Fall risk

- Cognition
- Mood
- Polypharmacy
- Social support
- Financial concerns
- Goals of care
- Advance care preferences

Additional components may also include evaluation of the following:

- Nutrition/weight change
- Urinary continence
- Sexual function
- Vision/hearing
- Dentition
- Living situation
- Spirituality

This section will focus on the core components of CGA. Although other aspects of the geriatric assessment are usually addressed during the CGA (eg, vision/hearing, nutrition), these components are discussed separately. (See "[Geriatric health maintenance](#)".)

Functional status — Functional status refers to the ability to perform activities necessary or desirable in daily life. Functional status is directly influenced by health conditions, particularly in the context of an elder's environment and social support network. Changes in functional status (eg, not being able to bathe independently) should prompt further diagnostic evaluation and intervention. Measurement of functional status can be valuable in monitoring response to treatment and can provide prognostic information that assists in long-term care planning.

Activities of daily living — An older adult's functional status can be assessed at three levels: basic activities of daily living (BADLs), instrumental or intermediate activities of daily living (IADLs), and advanced activities of daily living (AADLs).

BADLs refer to self-care tasks which include:

- Bathing
- Dressing
- Toileting
- Maintaining continence
- Grooming
- Feeding
- Transferring

IADLs refer to the ability to maintain an independent household which include:

- Shopping for groceries
- Driving or using public transportation
- Using the telephone
- Performing housework
- Doing home repair
- Preparing meals
- Doing laundry
- Taking medications
- Handling finances

Other possible IADLs that reflect the increased reliance on technology, which have not been validated, include:

- Ability to use a cellphone or smartphone
- Ability to use the internet
- Ability to keep a schedule of activities

AADLs vary considerably from individual to individual. These advanced activities include the ability to fulfill societal, community, and family roles as well as participate in recreational or occupational tasks.

Scales that measure functional status at each of these levels have been developed and validated. The [Vulnerable Elders Scale-13](#) (VES-13) is a 13-item screening tool that is based upon age, self-rated health, and the ability to perform functional and physical activities [7-9]. It identifies populations of community-dwelling elders at increased risk for functional decline or death over a five-year period (table 2). The VES-13 can be self-administered or administered by nonmedical personnel over the telephone or at an office visit in less than five minutes.

Questions that ask about specific BADL and IADL functions have also been incorporated into a variety of more generic, health-related quality-of-life instruments (eg, the Medical Outcomes Study Short-form and its shorter version, the SF-12; the PROMIS family of instruments) [8,10-12]. Two commonly used indices are the Katz index for ADLs (table 3) and the Lawton scale for IADLs (table 4). Some AADLs (eg, exercise and leisure time physical activity) can be ascertained by using standardized instruments. However, given the broad nature of AADLs, open-ended questions asking how one's day is spent might provide a better assessment of function in healthier older persons.

Adults over age 70 are more likely to have motor vehicle accidents as well as increased associated mortality. The patient's ability and safety to drive a car should also be evaluated in the functional assessment. (See ["Approach to the evaluation of older drivers"](#).)

Gait speed — In addition to measures of ADLs, gait speed alone predicts functional decline and early mortality in older adults [13]. Assessing gait speed in clinical practice may identify patients who need further evaluation, such as those at increased risk of falls. Additionally, assessing gait speed may help identify frail patients who might not benefit from treatment of chronic asymptomatic diseases such as hypertension. For example, elevated blood pressure in individuals age 65 and older was associated with increased mortality only in individuals with a walking speed ≥ 0.8 meters/second (measured over 6 meters or 20 feet) [14].

Falls/imbalance — Approximately one-third of community-dwelling persons age 65 years and one-half of those over 80 years of age fall each year. Patients who have fallen or have a gait or balance problem are at higher risk of having a subsequent fall and losing independence. An assessment of fall risk should be integrated into the history and physical examination of all geriatric patients ([algorithm 1](#)). (See "[Falls in older persons: Risk factors and patient evaluation](#)", [section on 'Falls risk assessment'](#) and "[Neurologic gait disorders of elderly people](#)".)

Cognition — The incidence of dementia increases with age, particularly among those over 85 years, yet many patients with cognitive impairment remain undiagnosed. The value of making an early diagnosis includes the possibility of uncovering treatable conditions. The evaluation of cognitive function can include a thorough history and brief cognition screens. If these raise suspicion for cognitive impairment, additional evaluation is indicated, which may include detailed mental status examination, neuropsychologic testing, tests to evaluate medical conditions that may contribute to cognitive impairment (eg, B12, thyroid-stimulating hormone [TSH]), depression assessment, and/or radiographic imaging (computed tomography [CT] or magnetic resonance imaging [MRI]). (See "[Evaluation of cognitive impairment and dementia](#)".)

Mood disorders — Depressive illness in the elder population is a serious health concern leading to unnecessary suffering, impaired functional status, increased mortality, and excessive use of health care resources. (See "[Diagnosis and management of late-life unipolar depression](#)".)

Late-life depression remains underdiagnosed and inadequately treated. Depression in elder adults may present atypically and may be difficult to assess in patients with cognitive impairment. A two-question screener is easily administered and likely to identify patients at risk if both questions are answered affirmatively [[15](#)]. The questions are:

- "During the past month, have you been bothered by feeling down, depressed, or hopeless?"
- "During the past month, have you been bothered by little interest or pleasure in doing things?"

This two-question screen is sensitive but not specific ([table 5](#)). Thus, a positive screen should be supplemented with seven additional questions to complete the Patient Health Questionnaire-9 (PHQ-9) [[16](#)]. The PHQ-9 has increasingly been used to detect and monitor depression symptoms among elder adults ([table 6](#)) [[17](#)]. The PHQ-9 provides a reliable and valid measure of depression severity.

A variety of other screens for depression are available and each has its advantages and disadvantages [[18](#)].

Polypharmacy — Older persons are often prescribed multiple medications by different health care providers, putting them at increased risk for drug-drug interactions and adverse drug events. The clinician should review the patient's medications at each visit. The best method of detecting potential problems with polypharmacy is to have patients bring in all of his/her medications (prescription and nonprescription) in their bottles. Discrepancies between what is documented in the medical record and what the patient is actually taking must be reconciled. As health systems have moved towards electronic health records and e-prescribing, the potential to detect potential medication errors and interactions has increased substantially. Although this can improve safety, record-generated messages about unimportant or rare interactions may lead to "reminder fatigue." (See "[Drug prescribing for older adults](#)" and "[Medical care in skilled nursing facilities \(SNFs\) in the United States](#)", [section on 'Prevention of adverse drug events'](#).)

Elder patients should also be asked about alternative medical therapy. As an example, herb use can be assessed by questioning: "What prescription medications, over the counter medicines, vitamins, herbs, or supplements do you use?" (See "[Overview of herbal medicine and dietary supplements](#)".)

Social and financial support — The existence of a strong social support network in an elder's life can frequently be the determining factor of whether the patient can remain at home or needs placement in an institution. A brief screen of social support includes taking a social history and determining who would be available to the elder to help if he or she becomes ill. Early identification of problems with social support can

help planning and timely development of resource referrals. For patients with functional impairment, the clinician should ascertain who the person has available to help with activities of daily living. (See '[Activities of daily living](#)' above.)

Caregivers should be screened periodically for symptoms of depression or caregiver burnout and, if present, referred for additional caregiving services, counseling, or support groups. Elder mistreatment should be considered in any geriatric assessment, particularly if the patient presents with contusions, burns, bite marks, genital or rectal trauma, pressure ulcers, or malnutrition with no clinical explanation. (See "[Elder mistreatment: Abuse, neglect, and financial exploitation](#)".)

The financial situation of a functionally impaired older adult is important to assess. Elders may qualify for state or local benefits, depending upon their income. Older patients occasionally have other benefits such as long-term care insurance or veteran's benefits that can help in paying for caregivers or prevent the need for institutionalization.

Goals of care — Most older adult patients who are appropriate for CGA have limited potential to return to fully healthy and independent lives. Hence, choices must be made about what outcomes are most important for them and their families. Goals of care often differ from advance care preferences that focus on future states of health that would be acceptable, determination of surrogates to make decisions, and medical treatments. Generally, advance directives are framed in the context of future deterioration in health status. (See '[Advance care preferences](#)' below.)

By contrast, a patient's goals of care are often positive (eg, regaining a previous health status, attending a future family event). Frequently, social (eg, living at home, maintaining social activities) and functional (eg, completing ADLs without help) goals assume priority over health-related goals (eg, survival) [19]. They are also patient-centric and individualized. For example, regaining independent ambulation after a hip fracture may be a goal for one patient whereas another might be content with use of a walker. Both short-term and longer-range goals should be considered and progress towards meeting these goals should be monitored, including reassessment if goals are not met within a specified time period. One approach that has been used in CGA is Goal Attainment Scaling [20,21]. However, clinicians can establish and monitor patient goals more informally by determining these in the course of clinical care and asking about them during subsequent visits.

Advance care preferences — Clinicians should begin discussions with all patients about preferences for specific treatments while the patient still has the cognitive capacity to make these decisions. These discussions should include preparation for in-the-moment decision-making [22], which includes choosing an appropriate decision-maker (ie, appointing a durable power of attorney, also known as a health care proxy, to serve as a surrogate in the event of personal incapacity), clarifying and articulating patients' values over time, and thinking about factors other than the patient's stated preferences in surrogate decision-making. As an example, patients who want to extend their life as long as possible might be asked about what should be done if the patient's health status changes and doctors recommend against further treatment, or if it becomes too hard for loved ones to keep them at home. Advance directives help guide therapy if a patient is unable to speak for him or herself and are vital to caring optimally for the geriatric population. (See "[Ethical issues in palliative care](#)".)

Tools have been developed to help providers elicit and document care preferences and promote shared end-of-life decision-making among seriously ill patients [5,6], including the Physician Orders for Life Sustaining Treatment (POLST) [23,24].

EFFICACY — Most meta-analyses have found that comprehensive geriatric assessment (CGA) leads to improved detection and documentation of geriatric problems [1,25-29]. However, the ability of CGA to improve outcomes (eg, decreased hospitalization, nursing home admission, and mortality) depends on specific CGA models and the settings where they have been implemented.

Several meta-analyses of randomized trials have evaluated five models of CGA [1,25-29]:

- Home geriatric assessment
- Acute geriatric care units
- Post-hospital discharge
- Outpatient consultation
- Inpatient consultation

Home geriatric assessment and acute geriatric care units have been shown to be consistently beneficial for several health outcomes. By contrast, the data are conflicting for post-hospital discharge, outpatient geriatric consultation, and inpatient geriatric consultation services.

Home assessment — Home geriatric assessment programs focus primarily on preventive rather than rehabilitative services. Although home assessment programs vary, most programs include a visiting nurse trained in geriatric care, as well as a physical therapist, social worker, psychologist, and specialty referrals when appropriate. In addition to home visits, telephone follow-up is routinely performed. Patients assessed at home are usually followed for at least one year. Although the number of frail older adults receiving home-based medical care is increasing, the majority of eligible homebound and home-limited older adults, particularly rural residents, have not received these services [30].

Multiple meta-analyses have found home assessments to be consistently effective in reducing functional decline as well as overall mortality [1,26-28]. As an example, a meta-analysis of 21 randomized trials found that multidimensional home visit programs were effective in reducing functional decline if a clinical examination was conducted (odds ratio [OR] 0.64, CI 0.48-0.87) and in reducing mortality in patients age ≤ 77 years old (OR 0.74, 95% CI 0.58-0.94) [28]. However, the home visits did not significantly prevent nursing home admissions (OR 0.86, CI 0.68-1.10). Like other meta-analyses for home assessments, this study was limited by heterogeneity across studies for all outcomes.

Acute geriatric care units — Several inpatient geriatric unit approaches have been developed in a variety of clinical settings. Within the US Department of Veterans Affairs Hospitals, these are usually referred to as Geriatric Evaluation and Management Units (GEMUs). In academic and private sector hospitals, they are usually labeled Acute Care of the Elderly (ACE) units. ACE units initially included structural modifications to promote mobility and simulate living conditions at home in preparation for a return to independence. More recently, however, ACE units are located on conventional hospital wards as designated geriatric units and tend to focus exclusively on processes of acute care rather than providing simulated living conditions and extended rehabilitation.

GEMUs are hospital wards that care for frail elderly patients through the multidisciplinary team approach. GEMUs have two main advantages over inpatient CGA consultation models. First, clinicians staffing the unit generally assume primary care of the patient, thus facilitating the implementation of recommendations. Second, the availability and experience of a dedicated team of providers (eg, nurses and therapists) increase the consistency and geriatric orientation of hospital care.

Numerous studies have compared the effect of inpatient CGA with usual care. A meta-analysis of 29 randomized trials involving nearly 14,000 participants found that patients who received CGA were more likely to be living at home (relative risk [RR] 1.06, 95% CI 1.01-1.10) and were less likely to be admitted to a nursing home up to a year after hospital admission (RR 0.80, 95% CI 0.72-0.89) [31]. There were no differences in dependence or cognitive status. CGA did not reduce the risk of death or the need for assistance with activities such as feeding or walking. This meta-analysis was limited by wide variability in interventions across studies. Moreover, due to length of stay (up to three months), such rehabilitative units are rarely available in the United States outside the Department of Veterans Affairs hospitals.

ACE units in acute care hospitals promote mobility and include processes to provide patient-centered care with nursing-initiated protocols. ACE units involve more intensive discharge planning and more detailed

education to improve medication compliance, in comparison with usual hospital care. In clinical trials, care in ACE units was associated with greater independence in ADLs at discharge, less frequent discharge to a nursing home, shorter and less expensive hospitalization [32,33], and reduced 30-day readmission rates [33], as well as higher satisfaction rates among patients, family members, clinicians, and nurses [34].

Due to the logistical barriers of having dedicated geriatric inpatient units (eg, unfilled beds when the census is light, overflow to other units when the unit is full), some programs have attempted to recreate the core elements of ACE units for hospitalized older persons who are not located on a single unit [35]. One matched cohort study indicated that benefits may include lower rates of adverse events, shorter hospital stays, and better satisfaction [36]. Whether these "virtual" units are as effective as ACE units is unknown; a 2011 meta-analysis suggests that they are not [37]. The lack of a consistent nursing staff that is trained in the care of older persons may diminish the effectiveness of this model.

Post-hospital discharge — Key elements of post-hospital discharge geriatric assessment include targeting criteria to identify vulnerable patients, a program of multidimensional assessment, comprehensive discharge planning, and home follow-up with nurses with specialized geriatrics training who visit the patients during the hospitalization and at least twice during the weeks following discharge. This intervention usually is initiated one to two days prior to hospital discharge. Similar to the home assessments discussed above, the post-discharge home visits are supplemented by telephone calls and additional visits by physical therapy, occupational therapy, social work, and/or home nursing services when indicated. (See '[Home assessment](#)' above.)

Studies of CGA have found inconsistent benefit for post-hospital discharge programs [1,38-41]. As an example, in a randomized trial of post-hospitalization CGA conducted in the home versus usual care, there was no difference between treatment and control arms in reducing functional decline, readmission rates, or mortality after 60 days [38]. In another randomized trial of comprehensive discharge planning with home follow-up versus usual care, there was no difference in functional status, post-discharge acute care visits, depression, or patient satisfaction after 24 weeks [39]. However, those randomly assigned to the intervention were less likely to be readmitted to the hospital compared with the control group (20 versus 37 percent, respectively). The intervention was also associated with a reduction in cost. Subsequent studies of similar discharge management programs with in-home follow-up have also found a reduction in readmission rates, for up to 12 months in some studies [41]. A systematic review found that many of the components of CGA were parts of care transition interventions that were effective in reducing rehospitalizations and emergency department visits [42].

CGA programs for patients discharged to home from the emergency department were found to be effective at reducing emergency department visits and hospital admission [40].

Outpatient consultation — Although meta-analyses have not shown benefit of outpatient CGA consultation [1,25], more complex CGA programs that address adherence to program recommendations and treat patients at higher risk of hospitalization have led to improved outcomes [43,44].

The first meta-analysis to evaluate CGA included four randomized trials and did not demonstrate benefit from outpatient CGA consultation in terms of hospital admission, nursing home placement, or physical/cognitive function [1]. However, one trial from this meta-analysis did not address whether recommendations from CGA were implemented and another trial included patients with poor prognoses, which may limit the generalizability of these data.

Some [43-46], but not all [47], of the subsequent randomized trials have shown some efficacy of outpatient CGA. Representative trials include:

- CGA coupled with an adherence intervention designed to empower the patient and prompt the primary care clinician to adhere to CGA recommendations was evaluated in a randomized trial [43]. Among community-dwelling older persons with functional disability, urinary incontinence, falls, or depressive

symptoms, those randomly assigned to CGA with adherence intervention had less functional decline, less fatigue, and better social functioning over a 15-month period compared with the control group.

- In a randomized trial evaluating CGA followed by six months of interdisciplinary primary care versus usual primary care in a population at risk for high health care utilization, those randomly assigned to treatment had reduced functional decline, depression, and use of home health services compared with the control group over 15 to 18 months following randomization [44].
- However, in a large, cluster-randomized trial of multidimensional geriatric assessment followed by either geriatric team management or the primary care clinician alone, there were no differences between the groups in hospitalization, admission to other institutions, and quality of life [47].

In a meta-analysis of nine randomized controlled trials (n = 3750) evaluating mortality, there was no benefit of outpatient CGA on survival (RR 0.95, 95% CI 0.82-1.12) [25]. Tests for heterogeneity showed consistency between trial data.

Specialized team management — Approaches to outpatient CGA have used some of the more successful components of older models and adapted them to programs within primary care practices:

- Geriatric Resources for Assessment and Care of Elders (GRACE) includes home-based CGA and long-term care management by a nurse practitioner and social worker who collaborate with the primary care clinician and a geriatrics interdisciplinary team. In a randomized trial of low-income elderly patients, those randomly assigned to the GRACE intervention had better health-related quality-of-life and fewer emergency department visits compared with those assigned to usual care [48]. Patients at high risk of hospitalization also had fewer admissions by the second year.
- Guided Care integrates an intensively trained nurse into primary care practices to provide geriatric assessment and chronic care management to high-risk patients. In a randomized trial of chronically ill older patients, those randomly assigned to Guided Care reported improved satisfaction rates and had less health care utilization with lower costs of care compared with those randomly assigned to usual care at eight months [49,50], but by 20 months of follow-up, the only significant reduction of health care was fewer episodes of home health care in the intervention group (OR 0.70, 95% CI 0.53-0.93). Among health maintenance organization (HMO) patients, the intervention also reduced the number of skilled nursing facility admissions (OR 0.53, CI 0.31-0.89) and days of hospitalization (OR 0.48, CI 0.28-0.84) [51].
- Practice redesign approaches (screening, structured visit notes, delegation to office staff, outreach to community resources) focus on specific geriatric conditions for assessment and management by clinicians or nurse practitioners. In two trials of patients in community-based practices, patients randomly assigned to practice-based interventions received better quality of care for falls and incontinence compared with those randomly assigned to usual care [45]. In a study within an academic geriatrics practice, this model of comanagement for five geriatric conditions resulted in improvements in quality of care for dementia, falls, and urinary incontinence when compared with a wait list control group; similar findings have been demonstrated in community-based practices [52,53].

Inpatient consultation — A meta-analysis of randomized trials of inpatient consultation found benefit for short-term (six- and eight-month) survival but no effect on functional status, readmission, or length of stay [54]. Two previous meta-analyses of inpatient consultation for CGA showed little benefit [1,55].

Many of the principles of inpatient CGA consultation have been incorporated into comanagement programs. As an example, comanagement with a geriatrician may reduce mortality, complications, delirium, and rehospitalization among patients with hip fracture on the surgical service [56-59]. A meta-analysis of surgical comanagement demonstrated decreased length of stay and suggested the possibility of decreased inpatient mortality [60]. In a second meta-analysis, CGA for persons with hip fracture provided as consultation or on an inpatient unit reduced the likelihood of being discharged to a setting where they would receive an increased

level of care such as an assisted-living or long-term care facility compared with discharge home; although not significant, the meta-analysis suggested a benefit on inpatient mortality [59].

New applications of comprehensive geriatric assessment — Principles and processes of CGA are increasingly being applied to subspecialty conditions, including cancer patients undergoing chemotherapy [61,62], considerations of surgery, or transcatheter aortic valve replacement (TAVR) for patients with aortic stenosis [63], vascular surgery [64], and postoperative mortality risk [65].

SUMMARY AND RECOMMENDATIONS

- Comprehensive geriatric assessment (CGA) is defined as a multidisciplinary diagnostic and treatment process that identifies medical, psychosocial, and functional capabilities of an older adult in order to develop a coordinated plan to maximize overall health with aging. CGA is based on the premise that a systematic evaluation of frail older persons by a team of health professionals may identify a variety of treatable health problems and lead to better health outcomes. (See ['Background'](#) above.)
- No standard criteria are available to readily identify patients who are likely to benefit from CGA. Specific criteria used by CGA programs to identify patients include:
 - Age
 - Medical comorbidities (eg, hip fracture, transcatheter aortic valve replacement [TAVR])
 - Psychosocial problems
 - Specific geriatric conditions such as functional disability
 - Previous or predicted high health care utilization
 - Consideration of change in living situation (eg, from independent living to assisted living, nursing home, or in-home caregivers)
- Community-dwelling older patients with functional disability, increased fall risk, cognitive decline, depression, or those at high risk for high health care utilization are appropriate candidates for CGA. Follow-up support is needed to ensure implementation of the findings of the CGA. (See ['Indications for referral'](#) above.)
- Inpatients admitted with fractures, failure to thrive, recurrent pneumonia, and pressure sores may benefit from hospitalization on a geriatric unit that provides CGA. Patients age ≥ 85 years who have geriatric conditions such as dementia or immobility should also be considered for care on these units. (See ['Indications for referral'](#) above.)
- The assessment team usually consists of a clinician, nurse, and social worker. Depending on the setting and patient's condition, the team may also include physical and occupational therapists, nutritionists, pharmacists, psychiatrists, psychologists, dentists, audiologists, podiatrists, and opticians. (See ['Assessment team'](#) above.)
- There are several components of CGA that should be evaluated, including (see ['Major components'](#) above):
 - Functional capacity
 - Fall risk
 - Cognition
 - Mood

- Polypharmacy
 - Nutrition/weight change
 - Urinary incontinence
 - Sexual function
 - Vision/hearing
 - Dentition
 - Living situation
 - Social support
 - Financial concerns
 - Goals of care
 - Spirituality
 - Advance care preferences
- Home geriatric assessment has been shown to be effective in improving functional status, preventing institutionalization, and reducing mortality. CGA performed in the hospital, especially in dedicated units, also has benefit on survival. Most programs of hospital discharge management with in-home follow-up have reduced readmission rates. However, studies of CGA have found inconsistent benefit for outpatient and inpatient geriatric consultation, except in the context of specific conditions (eg, hip fracture). (See ['Efficacy'](#) above.)

Use of UpToDate is subject to the [Subscription and License Agreement](#).

REFERENCES

1. Stuck AE, Siu AL, Wieland GD, et al. Comprehensive geriatric assessment: a meta-analysis of controlled trials. *Lancet* 1993; 342:1032.
2. Devons CA. Comprehensive geriatric assessment: making the most of the aging years. *Curr Opin Clin Nutr Metab Care* 2002; 5:19.
3. Garrison GM, Robelia PM, Pecina JL, Dawson NL. Comparing performance of 30-day readmission risk classifiers among hospitalized primary care patients. *J Eval Clin Pract* 2017; 23:524.
4. Elsayy B, Higgins KE. The geriatric assessment. *Am Fam Physician* 2011; 83:48.
5. http://www.geronet.ucla.edu/images/stories/docs/professionals/Geri_Pre-visit_Questionnaire.pdf (Accessed on November 14, 2011).
6. Reuben DB. Medical care for the final years of life: "When you're 83, it's not going to be 20 years". *JAMA* 2009; 302:2686.
7. Saliba D, Elliott M, Rubenstein LZ, et al. The Vulnerable Elders Survey: a tool for identifying vulnerable older people in the community. *J Am Geriatr Soc* 2001; 49:1691.
8. Min L, Yoon W, Mariano J, et al. The vulnerable elders-13 survey predicts 5-year functional decline and mortality outcomes in older ambulatory care patients. *J Am Geriatr Soc* 2009; 57:2070.
9. Min LC, Elliott MN, Wenger NS, Saliba D. Higher vulnerable elders survey scores predict death and functional decline in vulnerable older people. *J Am Geriatr Soc* 2006; 54:507.

10. Reuben DB, Borok GM, Wolde-Tsadik G, et al. A randomized trial of comprehensive geriatric assessment in the care of hospitalized patients. *N Engl J Med* 1995; 332:1345.
11. Ware JE, et al. SF12: an even shorter health survey. *Med Outcomes Trust Bull* 1996; 4:2.
12. <https://www.assessmentcenter.net/documents/InstrumentLibrary.pdf> (Accessed on August 05, 2015).
13. Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA* 2011; 305:50.
14. Odden MC, Peralta CA, Haan MN, Covinsky KE. Rethinking the association of high blood pressure with mortality in elderly adults: the impact of frailty. *Arch Intern Med* 2012; 172:1162.
15. Arroll B, Khin N, Kerse N. Screening for depression in primary care with two verbally asked questions: cross sectional study. *BMJ* 2003; 327:1144.
16. Rinaldi P, Mecocci P, Benedetti C, et al. Validation of the five-item geriatric depression scale in elderly subjects in three different settings. *J Am Geriatr Soc* 2003; 51:694.
17. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; 16:606.
18. Hoyl MT, Alessi CA, Harker JO, et al. Development and testing of a five-item version of the Geriatric Depression Scale. *J Am Geriatr Soc* 1999; 47:873.
19. Reuben DB, Tinetti ME. Goal-oriented patient care--an alternative health outcomes paradigm. *N Engl J Med* 2012; 366:777.
20. Rockwood K, Howlett S, Stadnyk K, et al. Responsiveness of goal attainment scaling in a randomized controlled trial of comprehensive geriatric assessment. *J Clin Epidemiol* 2003; 56:736.
21. Jennings LA, Ramirez KD, Hays RD, et al. Personalized Goal Attainment in Dementia Care: Measuring What Persons with Dementia and Their Caregivers Want. *J Am Geriatr Soc* 2018; 66:2120.
22. Sudore RL, Fried TR. Redefining the "planning" in advance care planning: preparing for end-of-life decision making. *Ann Intern Med* 2010; 153:256.
23. Austin CA, Mohottige D, Sudore RL, et al. Tools to Promote Shared Decision Making in Serious Illness: A Systematic Review. *JAMA Intern Med* 2015; 175:1213.
24. Oczkowski SJ, Chung HO, Hanvey L, et al. Communication Tools for End-of-Life Decision-Making in Ambulatory Care Settings: A Systematic Review and Meta-Analysis. *PLoS One* 2016; 11:e0150671.
25. Kuo HK, Scandrett KG, Dave J, Mitchell SL. The influence of outpatient comprehensive geriatric assessment on survival: a meta-analysis. *Arch Gerontol Geriatr* 2004; 39:245.
26. Stuck AE, Egger M, Hammer A, et al. Home visits to prevent nursing home admission and functional decline in elderly people: systematic review and meta-regression analysis. *JAMA* 2002; 287:1022.
27. Elkan R, Kendrick D, Dewey M, et al. Effectiveness of home based support for older people: systematic review and meta-analysis. *BMJ* 2001; 323:719.
28. Huss A, Stuck AE, Rubenstein LZ, et al. Multidimensional preventive home visit programs for community-dwelling older adults: a systematic review and meta-analysis of randomized controlled trials. *J Gerontol A Biol Sci Med Sci* 2008; 63:298.
29. Bachmann S, Finger C, Huss A, et al. Inpatient rehabilitation specifically designed for geriatric patients: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2010; 340:c1718.
30. Yao NA, Ritchie C, Cornwell T, Leff B. Use of Home-Based Medical Care and Disparities. *J Am Geriatr Soc* 2018; 66:1716.
31. Ellis G, Gardner M, Tsiachristas A, et al. Comprehensive geriatric assessment for older adults admitted to hospital. *Cochrane Database Syst Rev* 2017; 9:CD006211.
32. Landefeld CS, Palmer RM, Kresevic DM, et al. A randomized trial of care in a hospital medical unit especially designed to improve the functional outcomes of acutely ill older patients. *N Engl J Med* 1995; 332:1338.

33. Flood KL, MacLennan PA, McGrew D, et al. Effects of an acute care for elders unit on costs and 30-day readmissions. *JAMA Intern Med* 2013; 173:981.
34. Counsell SR, Holder CM, Liebenauer LL, et al. Effects of a multicomponent intervention on functional outcomes and process of care in hospitalized older patients: a randomized controlled trial of Acute Care for Elders (ACE) in a community hospital. *J Am Geriatr Soc* 2000; 48:1572.
35. Arbaje AI, Maron DD, Yu Q, et al. The geriatric floating interdisciplinary transition team. *J Am Geriatr Soc* 2010; 58:364.
36. Hung WW, Ross JS, Farber J, Siu AL. Evaluation of the Mobile Acute Care of the Elderly (MACE) service. *JAMA Intern Med* 2013; 173:990.
37. Ellis G, Whitehead MA, Robinson D, et al. Comprehensive geriatric assessment for older adults admitted to hospital: meta-analysis of randomised controlled trials. *BMJ* 2011; 343:d6553.
38. Siu AL, Kravitz RL, Keeler E, et al. Postdischarge geriatric assessment of hospitalized frail elderly patients. *Arch Intern Med* 1996; 156:76.
39. Naylor MD, Brooten D, Campbell R, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA* 1999; 281:613.
40. Caplan GA, Williams AJ, Daly B, Abraham K. A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department--the DEED II study. *J Am Geriatr Soc* 2004; 52:1417.
41. Naylor MD, Aiken LH, Kurtzman ET, et al. The care span: The importance of transitional care in achieving health reform. *Health Aff (Millwood)* 2011; 30:746.
42. Hesselink G, Schoonhoven L, Barach P, et al. Improving patient handovers from hospital to primary care: a systematic review. *Ann Intern Med* 2012; 157:417.
43. Reuben DB, Frank JC, Hirsch SH, et al. A randomized clinical trial of outpatient comprehensive geriatric assessment coupled with an intervention to increase adherence to recommendations. *J Am Geriatr Soc* 1999; 47:269.
44. Boulton C, Boulton LB, Morishita L, et al. A randomized clinical trial of outpatient geriatric evaluation and management. *J Am Geriatr Soc* 2001; 49:351.
45. Wenger NS, Roth CP, Shekelle PG, et al. A practice-based intervention to improve primary care for falls, urinary incontinence, and dementia. *J Am Geriatr Soc* 2009; 57:547.
46. Burns R, Nichols LO, Martindale-Adams J, Graney MJ. Interdisciplinary geriatric primary care evaluation and management: two-year outcomes. *J Am Geriatr Soc* 2000; 48:8.
47. Fletcher AE, Price GM, Ng ES, et al. Population-based multidimensional assessment of older people in UK general practice: a cluster-randomised factorial trial. *Lancet* 2004; 364:1667.
48. Counsell SR, Callahan CM, Clark DO, et al. Geriatric care management for low-income seniors: a randomized controlled trial. *JAMA* 2007; 298:2623.
49. Boulton C, Reider L, Frey K, et al. Early effects of "Guided Care" on the quality of health care for multimorbid older persons: a cluster-randomized controlled trial. *J Gerontol A Biol Sci Med Sci* 2008; 63:321.
50. Leff B, Reider L, Frick KD, et al. Guided care and the cost of complex healthcare: a preliminary report. *Am J Manag Care* 2009; 15:555.
51. Boulton C, Reider L, Leff B, et al. The effect of guided care teams on the use of health services: results from a cluster-randomized controlled trial. *Arch Intern Med* 2011; 171:460.
52. Ganz DA, Koretz BK, Bail JK, et al. Nurse practitioner comanagement for patients in an academic geriatric practice. *Am J Manag Care* 2010; 16:e343.
53. Reuben DB, Ganz DA, Roth CP, et al. Effect of nurse practitioner comanagement on the care of geriatric conditions. *J Am Geriatr Soc* 2013; 61:857.

54. Deschodt M, Flamaing J, Haentjens P, et al. Impact of geriatric consultation teams on clinical outcome in acute hospitals: a systematic review and meta-analysis. *BMC Med* 2013; 11:48.
55. Ellis G, Whitehead MA, O'Neill D, et al. Comprehensive geriatric assessment for older adults admitted to hospital. *Cochrane Database Syst Rev* 2011; :CD006211.
56. Shields L, Henderson V, Caslake R. Comprehensive Geriatric Assessment for Prevention of Delirium After Hip Fracture: A Systematic Review of Randomized Controlled Trials. *J Am Geriatr Soc* 2017; 65:1559.
57. Fisher AA, Davis MW, Rubenach SE, et al. Outcomes for older patients with hip fractures: the impact of orthopedic and geriatric medicine cocare. *J Orthop Trauma* 2006; 20:172.
58. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc* 2001; 49:516.
59. Eamer G, Taheri A, Chen SS, et al. Comprehensive geriatric assessment for older people admitted to a surgical service. *Cochrane Database Syst Rev* 2018; 1:CD012485.
60. Van Grootven B, Flamaing J, Dierckx de Casterlé B, et al. Effectiveness of in-hospital geriatric co-management: a systematic review and meta-analysis. *Age Ageing* 2017; 46:903.
61. Kalsi T, Babic-Illman G, Ross PJ, et al. The impact of comprehensive geriatric assessment interventions on tolerance to chemotherapy in older people. *Br J Cancer* 2015; 112:1435.
62. Corre R, Greillier L, Le Caër H, et al. Use of a Comprehensive Geriatric Assessment for the Management of Elderly Patients With Advanced Non-Small-Cell Lung Cancer: The Phase III Randomized ESOGIA-GFPC-GECP 08-02 Study. *J Clin Oncol* 2016; 34:1476.
63. Boureau AS, Trochu JN, Colliard C, et al. Determinants in treatment decision-making in older patients with symptomatic severe aortic stenosis. *Maturitas* 2015; 82:128.
64. Partridge JS, Harari D, Martin FC, et al. Randomized clinical trial of comprehensive geriatric assessment and optimization in vascular surgery. *Br J Surg* 2017; 104:679.
65. Kim SW, Han HS, Jung HW, et al. Multidimensional frailty score for the prediction of postoperative mortality risk. *JAMA Surg* 2014; 149:633.

GRAPHICS

Questions and simple tests for general screening assessment of frail older patients*

	Question	Indicator (Scoring applies to individual domains)	Alternative
Functional status			
Activities of daily living (ADLs)	Bathing, dressing, toileting, transferring, maintaining continence, feeding	Able to complete without assistance; able but with difficulty; unable to complete without assistance	
Instrumental ADLs (IADLs)	Using the telephone, shopping, preparing meals, housekeeping, doing laundry, using public transportation or driving, taking medication, handling finances	Able to complete without assistance; unable to complete without assistance	
Visual impairment	Do you have difficulty driving, watching television, reading, or doing any of your daily activities because of your eyesight, even while wearing glasses? ^[1]	Yes indicates positive screen	Snellen eye chart
Hearing impairment [¶]	Is your age older than 70 years?	1 point	Alternative is Audioscope ^[2]
	Are you of male gender?	1 point	
	Do you have 12 or fewer years of education?	1 point	
	Did you ever see a doctor about trouble hearing?	2 points	
	Without a hearing aid, can you usually hear and understand what a person says without seeing his face if that person whispers to you from across the room?	If no, 1 point	
	Without a hearing aid, can you usually hear and understand what a person says without seeing his face if that person talks in a normal voice to you from across the room?	If no, 2 points	
		≥3 points: positive screen	
Urinary incontinence ^Δ	Have you had urinary incontinence (lose your urine) that is bothersome enough that you would like to know how it could be treated?	Yes indicates positive screen	
Malnutrition	Have you lost any weight in the last year? ^[3]	Loss of at least 5 percent of usual body weight in last year indicates positive screen ^[3]	
Gait, balance, falls ^Δ	Have you fallen two or more times in the past 12 months?	Any yes response indicates positive screen	
	Have you fallen and hurt yourself since your last doctor's visit?		
	Have you been afraid of falling because of balance or walking problems?		

Depression [◇]	Over the past two weeks, how often have you been bothered by:	Response score for each:	
	Little interest or pleasure in doing things?	0: not at all 1: several days 2: more than half the days 3: nearly every day	
	Feeling down, depressed, or hopeless?	Total ≥3, positive screen	
Cognitive problems	Three-item recall ^[4]	<2 items recalled indicates positive screen ^[4]	
	Clock-drawing test ^[5]	Any of the following errors indicate positive screen: wrong time, no hands, missing numbers, number substitutions, repetition, refusal ^[5]	
Environmental problems	Home safety checklists ^[6]		

* All except the Snellen eye chart, Audioscope, and evaluation for cognitive problems can be assessed by self-report using questionnaire.

¶ Questions and response indicators are from the National Health and Nutrition Examination Survey (NHANES) battery.^[7]

Δ Questions and response indicators are from the ACOVE-2 Screener.^[8]

◇ Questions and response indicators are from the Patient Health Questionnaire-2.^[9]

References:

1. Moore AA, Siu AL. Screening for common problems in ambulatory elderly. *Am J Med* 1996; 100:438.
2. Yueh B, Shapiro N, MacLean CH, Shekelle PG. Screening and management of adult hearing loss in primary care. *JAMA* 2003; 289:1976.
3. Wallace JI, Schwartz RS, LaCroix AZ, Uhlmann RF, Pearlman RA. Involuntary weight loss in older outpatients. *J Am Geriatr Soc* 1995; 43:329.
4. Siu AL. Screening for dementia and investigating its causes. *Ann Intern Med* 1991; 115:122.
5. Lessig MC, Scanlan JM, Nazemi H, Borson S. Time that tells: critical clockdrawing errors for dementia screening. *Int Psychogeriatr* 2008; 20:459.
6. Department of Health and Human Services; Centers for Disease Control and Prevention; National Center for Injury Prevention and Control. Home safety checklists. <http://www.cdc.gov/ncjpc/falls/FallPrev4.pdf>. Accessed June 9, 2009.
7. Reuben DB, Walsh K, Moore AA, et al. Hearing loss in community-dwelling older persons. *J Am Geriatr Soc* 1998; 46:1008.
8. Wenger NS, Roth CP, Shekelle PG, et al. Practice-based intervention to improve primary care for falls, urinary incontinence and dementia. *J Am Geriatr Soc* 2009; 57:547.
9. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2. *Med Care* 2003; 41:1284.

Reproduced with permission from: Rueben DB. Medical care for the final years of life: "When you're 83, it's not going to be 20 years." *JAMA* 2009; 302:2686. Copyright © 2009 American Medical Association. All rights reserved.

The Vulnerable Elders Survey (VES) 13 scale

Domain	Score
Age	
75-85	1
>85	3
Self-rated health	
Good, very good, and excellent	0
Fair and poor	1
Activities of daily living (ADLs)/instrumental ADLs (IADLs)	
Needs assistance with	
Bathing or showering	1
Shopping	1
Money management	1
Transfer	1
Light housework	1
Difficulty in special activities	
Kneeling, bending, and stooping	1
Performance of housework (example: scrubbing the floor)	1
Reaching out and lifting upper extremities above the shoulder	1
Lifting and carrying 10 pounds	1
Walking 0.25 miles	1
Writing or handling and grasping small objects	1

Score ≥ 3 : Vulnerable elderly

Data from: Saliba D, Elliott M, Rubenstein LZ, et al. The Vulnerable Elders Survey: a tool for identifying vulnerable older people in the community. *J Am Geriatr Soc* 2001; 49:1691.

Graphic 73154 Version 3.0

Katz index of independence in activities of daily living

Activities	Independence	Dependence
Points (1 or 0)	Points (1) NO supervision, direction, or personal assistance	Points (0) WITH supervision, direction, personal assistance, or total care
Bathing POINTS: _____	(1 point) Bathes self completely or needs help in bathing only a single part of the body, such as the back, genital area, or disabled extremity.	(0 points) Needs help with bathing more than one part of the body, getting in or out of the tub or shower. Requires total bathing.
Dressing POINTS: _____	(1 point) Gets clothes from closets and drawers and puts on clothes and outer garments complete with fasteners. May have help tying shoes.	(0 points) Needs help with dressing self or needs to be completely dressed.
Toileting POINTS: _____	(1 point) Goes to toilet, gets on and off, arranges clothes, cleans genital area without help.	(0 points) Needs help transferring to the toilet and cleaning self or uses bedpan or commode.
Transferring POINTS: _____	(1 point) Moves in and out of bed or chair unassisted. Mechanical transferring aides are acceptable.	(0 points) Needs help in moving from bed to chair or requires a complete transfer.
Continence POINTS: _____	(1 point) Exercises complete self-control over urination and defecation.	(0 points) Is partially or totally incontinent of bowel or bladder.
Feeding POINTS: _____	(1 point) Gets food from plate into mouth without help. Preparation of food may be done by another person.	(0 points) Needs partial or total help with feeding or requires parenteral feeding.
Total points: _____		

6 points: High (patient independent).

0 points: Low (patient very dependent).

Reproduced with permission from: Katz S, Down TD, Cash HR, Grotz RC. Progress in the development of the index of ADL. Gerontologist 1970, 10:20. Copyright © 1970 Oxford University Press.

Graphic 66451 Version 11.0


The Lawton instrumental activities of daily living scale

Activities	Points	Activities	Points
Ability to use telephone		Laundry	
1. Operates telephone on own initiative; looks up and dials numbers	1	1. Does personal laundry completely	1
2. Dials a few well-known numbers	1	2. Launders small items, rinses socks, stockings, etc	1
3. Answers telephone, but does not dial	1	3. All laundry must be done by others	0
4. Does not use telephone at all	0	Mode of transportation	
Shopping		1. Travels independently on public transportation or drives own car	1
1. Takes care of all shopping needs independently	1	2. Arranges own travel via taxi, but does not otherwise use public transportation	1
2. Shops independently for small purchases	0	3. Travels on public transportation when assisted or accompanied by another	1
3. Needs to be accompanied on any shopping trip	0	4. Travel limited to taxi or automobile with assistance of another	0
4. Completely unable to shop	0	5. Does not travel at all	0
Food preparation		Responsibility for own medications	
1. Plans, prepares, and serves adequate meals independently	1	1. Is responsible for taking medication in correct doses at correct time	1
2. Prepares adequate meals if supplied with ingredients	0	2. Takes responsibility if medication is prepared in advance in separate doses	0
3. Heats and serves prepared meals or prepares meals, but does not maintain adequate diet	0	3. Is not capable of dispensing own medication	0
4. Needs to have meals prepared and served	0	Ability to handle finances	
Housekeeping		1. Manages financial matters independently (budgets, writes checks, pays rent and bills, goes to bank); collects and keeps track of income	1
1. Maintains house alone with occasion assistance (heavy work)	1	2. Manages day-to-day purchases, but needs help with banking, major purchases, etc	1
2. Performs light daily tasks such as dishwashing, bed making	1	3. Incapable of handling money	0
3. Performs light daily tasks, but cannot maintain acceptable level of cleanliness	1		
4. Needs help with all home maintenance tasks	1		
5. Does not participate in any housekeeping tasks	0		

Scoring: For each category, circle the item description that most closely resembles the client's highest functional level (either 0 or 1).

A summary score ranges from 0 (dependent, requires significant assistance to live in the community) to 8 (independent, no assistance required to maintain self in community).

Prevention falls algorithm



Reproduced with permission from: *The Prevention of Falls in Older Persons: Clinical Practice Guideline* (<http://www.medcats.com/FALLS/frameset.htm>) from the American Geriatrics Society. For more information visit the AGS online at www.americangeriatrics.org.

Screening instruments for late-life depression for use in primary care

	Sensitivity percent	Specificity percent	Inpatient	Outpatient	Physically ill	Cognitively impaired
Two-question screen	97	67	Unknown	Yes	Unknown	No
Geriatric Depression Scale (5-item)	94	81	Yes	Yes	Yes	Unknown
Patient Health Questionnaire-9 (9-item)	88	88	Unknown	Yes	Yes	Unknown
Cornell Scale for Depression in Dementia (19-item)	90	75	Yes	Yes	Unknown	Yes
Center for Epidemiologic Studies - Depression Scale (20-item)	93	73	No	Yes	Unknown	No

Graphic 59906 Version 2.0

PHQ-9 depression questionnaire

Name:	Date:			
Over the last two weeks, how often have you been bothered by any of the following problems?	Not at all	Several days	More than half the days	Nearly every day
Little interest or pleasure in doing things	0	1	2	3
Feeling down, depressed, or hopeless	0	1	2	3
Trouble falling or staying asleep, or sleeping too much	0	1	2	3
Feeling tired or having little energy	0	1	2	3
Poor appetite or overeating	0	1	2	3
Feeling bad about yourself, or that you are a failure, or that you have let yourself or your family down	0	1	2	3
Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
Moving or speaking so slowly that other people could have noticed? Or the opposite, being so fidgety or restless that you have been moving around a lot more than usual.	0	1	2	3
Thoughts that you would be better off dead, or of hurting yourself in some way	0	1	2	3
Total ____ =	____	+ ____	+ ____	+ ____
PHQ-9 score ≥10: Likely major depression				
Depression score ranges:				
5 to 9: mild				
10 to 14: moderate				
15 to 19: moderately severe				
≥20: severe				
If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?	Not difficult at all ____	Somewhat difficult ____	Very difficult ____	Extremely difficult ____

PHQ: Patient Health Questionnaire.

Developed by Drs. Robert L Spitzer, Janet BW Williams, Kurt Kroenke, and colleagues, with an educational grant from Pfizer, Inc. No permission required to reproduce, translate, display or distribute.

Graphic 59307 Version 10.0

Contributor Disclosures

Katherine T Ward, MD Nothing to disclose **David B Reuben, MD** Nothing to disclose **Kenneth E Schmader, MD** Grant/Research/Clinical Trial Support: Merck [Herpes zoster (Zoster vaccine)]; GlaxoSmithKline [Herpes zoster (Zoster vaccine)]. **Jane Givens, MD** Consultant/Advisory Boards (Partner): CVS Health/CVS Omnicare [Pharmaceutical management of formulary decision-making].

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

[Conflict of interest policy.](#)